

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199914180 B2
(10) Patent No. 760010

(54) Title
DNA19355 polypeptide, a tumor necrosis factor homolog

(51)⁶ International Patent Classification(s)
C12N 015/28 C07K 016/24
A61K 038/19 C07K 019/00
C07K 014/525

(21) Application No: 199914180 (22) Application Date: 1998 .11 .18

(87) WIPO No: WO99/25834

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/065635	1997 .11 .18	US
60/069661	1997 .12 .12	US

(43) Publication Date : 1999 .06 .07
(43) Publication Journal Date : 1999 .08 .05
(44) Accepted Journal Date : 2003 .05 .08

(71) Applicant(s)
Genentech, Inc.

(72) Inventor(s)
Avi J Ashkenazi; Austin L. Gurney; Scot A.
Marsters ; Robert Pitti; Kevin P. Baker; Paul J Godowski;
Melanie R Mark

(74) Agent/Attorney
GRIFFITH HACK.GPO Box 1285K.MELBOURNE VIC 3001

(56) Related Art
AU 68478/96

60065635
60069661
PCT/97

14180/99

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 15/28, C07K 14/525, A61K 38/19, C07K 19/00, 16/24		A1	(11) International Publication Number: WO 99/25834
			(43) International Publication Date: 27 May 1999 (27.05.99)
(21) International Application Number: PCT/US98/24621		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TI, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 18 November 1998 (18.11.98)		Published With international search report.	
(30) Priority Data: 60/065,635 18 November 1997 (18.11.97) US 60/069,661 12 December 1997 (12.12.97) US			
(71) Applicant: GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080 (US).			
(72) Inventors: ASHKENAZI, Avi, J.; 1456 Terrytown Street, San Mateo, CA 94402 (US). GURNEY, Austin, L.; 1 Debbie Lane, Belmont, CA 94002 (US). MARSTERS, Scot, A.; 990 Cherry Street, San Carlos, CA 94070 (US). PITTI, Robert; 1110 Liberty Street, El Cerrito, CA 94530 (US). BAKER, Kevin, P.; 1115 South Grant Street, San Mateo, CA 94401 (US). GODOWSKI, Paul, J.; 2627 Easton Drive, Burlingame, CA 94010 (US). MARK, Melanie, R.; 958B Chula Vista Avenue, Burlingame, CA 94010 (US).			
(74) Agent: DAIGNAULT, Ronald, A.; Merchant, Gould, Smith, Edell, Welter & Schmidt, P.A., 3100 Norwest Center, 90 South Seventh Street, Minneapolis, MN 55402-4131 (US).			
(54) Title: DNA19355 POLYPEPTIDE, A TUMOR NECROSIS FACTOR HOMOLOG			
(57) Abstract A tumor necrosis factor homolog, identified as DNA19355, is provided. DNA19355 polypeptide has apoptotic activity in mammalian cancer cells and may be involved in proinflammatory responses. Nucleic acid molecules encoding DNA19355, chimeric molecules and antibodies to DNA19355 are also provided.			

DNA19355 POLYPEPTIDE, A TUMOUR NECROSIS FACTOR HOMOLOG

FIELD OF THE INVENTION

The present invention relates generally to the
5 identification and isolation of novel DNA and to the
recombinant production of novel polypeptides, designated
herein as "DNA19355".

BACKGROUND OF THE INVENTION

10 All references, including any patents or patent
applications, cited in this specification are hereby
incorporated by reference. No admission is made that any
reference constitutes prior art. The discussion of the
references states what their authors assert, and the
15 applicants reserve the right to challenge the accuracy and
pertinency of the cited documents. It will be clearly
understood that, although a number of prior art
publications are referred to herein, this reference does
not constitute an admission that any of these documents
20 forms part of the common general knowledge in the art, in
Australia or in any other country.

Control of cell numbers in mammals is believed to
be determined, in part, by a balance between cell
proliferation and cell death. One form of cell death,
25 sometimes referred to as necrotic cell death, is typically
characterised as a pathologic form of cell death resulting
from some trauma or cellular injury. In contrast, there
is another, "physiological" form of cell death which
usually proceeds in an orderly or controlled manner. This
30 orderly or controlled form of cell death is often referred
to as "apoptosis" [see, e.g., Barr et al., Bio/Technology,
12:487-493 (1994); Steller et al., Science, 267:1445-1449
(1995)]. Apoptotic cell death naturally occurs in many
physiological processes, including embryonic development
35 and clonal selection in the immune system [Itoh et al.,
Cell, 66:233-243 (1991)]. Decreased levels of apoptotic
cell death have been associated with a variety of



pathological conditions, including cancer, lupus, and
herpes virus infection [Thompson, Science, 267:1456-1462
(1995)]. Increased levels of apoptotic cell death may be
associated with a variety of other pathological
5 conditions, including AIDS, Alzheimer's disease,
Parkinson's disease, amyotrophic lateral sclerosis,
multiple sclerosis, retinitis pigmentosa, cerebellar
degeneration, aplastic anemia, myocardial infarction,
stroke, reperfusion injury, and toxin-induced liver
10 disease [see, Thompson, supra].

3
4

5
6
7
8



Apoptotic cell death is typically accompanied by one or more characteristic morphological and biochemical changes in cells, such as condensation of cytoplasm, loss of plasma membrane microvilli, segmentation of the nucleus, degradation of chromosomal DNA or loss of mitochondrial function. A variety of extrinsic and intrinsic signals are believed to trigger or induce such morphological and biochemical cellular changes [Raff, Nature, 356:397-400 (1992); Steller, supra; Sachs et al., Blood, 82:15 (1993)]. For instance, they can be triggered by hormonal stimuli, such as glucocorticoid hormones for immature thymocytes, as well as withdrawal of certain growth factors [Watanabe-Fukunaga et al., Nature, 356:314-317 (1992)]. Also, some identified oncogenes such as *myc*, *rel*, and *ELA*, and tumor suppressors, like *p53*, have been reported to have a role in inducing apoptosis. Certain chemotherapy drugs and some forms of radiation have likewise been observed to have apoptosis-inducing activity [Thompson, supra].

Various molecules, such as tumor necrosis factor- α ("TNF- α "), tumor necrosis factor- β ("TNF- β " or "lymphotoxin- α "), lymphotoxin- β ("LT- β "), CD30 ligand, CD27 ligand, CD40 ligand, OX-40 ligand, 4-1BB ligand, Apo-1 ligand (also referred to as Fas ligand or CD95 ligand), and Apo-2 ligand (also referred to as TRAIL) have been identified as members of the tumor necrosis factor ("TNF") family of cytokines [See, e.g., Gruss and Dower, Blood, 85:3378-3404 (1995); Pitti et al., J. Biol. Chem., 271:12687-12690 (1996); Wiley et al., Immunity, 3:673-682 (1995); Browning et al., Cell, 72:847-856 (1993); Armitage et al. Nature, 357:80-82 (1992)]. Among these molecules, TNF- α , TNF- β , CD30 ligand, 4-1BB ligand, Apo-1 ligand, and Apo-2 ligand (TRAIL) have been reported to be involved in apoptotic cell death. Both TNF- α and TNF- β have been reported to induce apoptotic death in susceptible tumor cells [Schmid et al., Proc. Natl. Acad. Sci., 83:1881 (1986); Dealtry et al., Eur. J. Immunol., 17:689 (1987)]. Zheng et al. have reported that TNF- α is involved in post-stimulation apoptosis of CD8-positive T cells [Zheng et al., Nature, 377:348-351 (1995)]. Other investigators have reported that CD30 ligand may be involved in deletion of self-reactive T cells in the thymus [Amakawa et al., Cold Spring Harbor

Laboratory Symposium on Programmed Cell Death, Abstr. No. 10, (1995)).

5 Mutations in the mouse Fas/Apo-1 receptor or ligand genes (called *lpr* and *gld*, respectively) have been associated with some autoimmune disorders, indicating that Apo-1 ligand may play a role in regulating the clonal deletion of self-reactive lymphocytes in the periphery [Krammer et al., Curr. Op. Immunol., 6:279-289 (1994); Nagata et al., Science, 267:1449-1456 (1995)]. Apo-1 ligand is also reported to induce post-stimulation apoptosis in CD4-
10 positive T lymphocytes and in B lymphocytes, and may be involved in the elimination of activated lymphocytes when their function is no longer needed [Krammer et al., supra; Nagata et al., supra]. Agonist mouse monoclonal antibodies specifically binding to the Apo-1 receptor have been reported to exhibit cell killing activity that
15 is comparable to or similar to that of TNF- α [Yonehara et al., J. Exp. Med., 169:1747-1756 (1989)].

Induction of various cellular responses mediated by such TNF family cytokines is believed to be initiated by their binding to specific cell receptors. Two distinct TNF receptors of
20 approximately 55-kDa (TNFR1) and 75-kDa (TNFR2) have been identified [Hohman et al., J. Biol. Chem., 264:14927-14934 (1989); Brockhaus et al., Proc. Natl. Acad. Sci., 87:3127-3131 (1990); EP 417,563, published March 20, 1991] and human and mouse cDNAs corresponding to both receptor types have been isolated and characterized [Loetscher
25 et al., Cell, 61:351 (1990); Schall et al., Cell, 61:361 (1990); Smith et al., Science, 248:1019-1023 (1990); Lewis et al., Proc. Natl. Acad. Sci., 88:2830-2834 (1991); Goodwin et al., Mol. Cell. Biol., 11:3020-3026 (1991)]. Extensive polymorphisms have been associated with both TNF receptor genes [see, e.g., Takao et al.,
30 Immunogenetics, 37:199-203 (1993)]. Both TNFRs share the typical structure of cell surface receptors including extracellular, transmembrane and intracellular regions. The extracellular portions of both receptors are found naturally also as soluble TNF-binding proteins [Nophar, Y. et al., EMBO J., 9:3269 (1990); and Kohno, T.
35 et al., Proc. Natl. Acad. Sci. U.S.A., 87:8331 (1990)]. More recently, the cloning of recombinant soluble TNF receptors was

reported by Hale et al. [J. Cell. Biochem. Supplement 15F, 1991, p. 113 (P424)].

The extracellular portion of type 1 and type 2 TNFRs (TNFR1 and TNFR2) contains a repetitive amino acid sequence pattern of four cysteine-rich domains (CRDs) designated 1 through 4, starting from the NH₂-terminus. Each CRD is about 40 amino acids long and contains 4 to 6 cysteine residues at positions which are well conserved [Schall et al., supra; Loetscher et al., supra; Smith et al., supra; Nophar et al., supra; Kohno et al., supra]. In TNFR1, the approximate boundaries of the four CRDs are as follows: CRD1- amino acids 14 to about 53; CRD2- amino acids from about 54 to about 97; CRD3- amino acids from about 98 to about 138; CRD4- amino acids from about 139 to about 167. In TNFR2, CRD1 includes amino acids 17 to about 54; CRD2- amino acids from about 55 to about 97; CRD3- amino acids from about 98 to about 140; and CRD4- amino acids from about 141 to about 179 [Banner et al., Cell, 73:431-435 (1993)]. The potential role of the CRDs in ligand binding is also described by Banner et al., supra.

A similar repetitive pattern of CRDs exists in several other cell-surface proteins, including the p75 nerve growth factor receptor (NGFR) [Johnson et al., Cell, 47:545 (1986); Radeke et al., Nature, 325:593 (1987)], the B cell antigen CD40 [Stamenkovic et al., EMBO J., 8:1403 (1989)], the T cell antigen OX40 [Mallet et al., EMBO J., 9:1063 (1990)] and the Fas antigen [Yonehara et al., supra and Itoh et al., Cell, 66:233-243 (1991)]. CRDs are also found in the soluble TNFR (sTNFR)-like T2 proteins of the Shope and myxoma poxviruses [Upton et al., Virology, 160:20-29 (1987); Smith et al., Biochem. Biophys. Res. Commun., 176:335 (1991); Upton et al., Virology, 184:370 (1991)]. Optimal alignment of these sequences indicates that the positions of the cysteine residues are well conserved. These receptors are sometimes collectively referred to as members of the TNF/NGF receptor superfamily. Recent studies on p75NGFR showed that the deletion of CRD1 [Welcher, A.A. et al., Proc. Natl. Acad. Sci. USA, 88:159-163 (1991)] or a 5-amino acid insertion in this domain [Yan, H. and Chao, M.V., J. Biol. Chem., 266:12099-12104 (1991)] had little or no effect on NGF binding [Yan, H. and Chao, M.V., supra]. p75 NGFR contains a proline-rich stretch of about 60 amino acids, between its CRD4 and

transmembrane region, which is not involved in NGF binding [Peetre, C. et al., Eur. J. Hematol., 41:414-419 (1988); Seckinger, P. et al., J. Biol. Chem., 264:11966-11973 (1989); Yan, H. and Chao, M.V., supra]. A similar proline-rich region is found in TNFR2 but not in TNFR1.

Itoh et al. disclose that the Apo-1 receptor can signal an apoptotic cell death similar to that signaled by the 55-kDa TNFR1 [Itoh et al., supra]. Expression of the Apo-1 antigen has also been reported to be down-regulated along with that of TNFR1 when cells are treated with either TNF- α or anti-Apo-1 mouse monoclonal antibody [Krammer et al., supra; Nagata et al., supra]. Accordingly, some investigators have hypothesized that cell lines that co-express both Apo-1 and TNFR1 receptors may mediate cell killing through common signaling pathways [Id.].

The TNF family ligands identified to date, with the exception of lymphotoxin- α , are type II transmembrane proteins, whose C-terminus is extracellular. In contrast, most receptors in the TNF receptor (TNFR) family identified to date are type I transmembrane proteins. In both the TNF ligand and receptor families, however, homology identified between family members has been found mainly in the extracellular domain ("ECD"). Several of the TNF family cytokines, including TNF- α , Apo-1 ligand and CD40 ligand, are cleaved proteolytically at the cell surface; the resulting protein in each case typically forms a homotrimeric molecule that functions as a soluble cytokine. TNF receptor family proteins are also usually cleaved proteolytically to release soluble receptor ECDs that can function as inhibitors of the cognate cytokines.

Recently, other members of the TNFR family have been identified. Such newly identified members of the TNFR family include CAR1, HVEM and osteoprotegerin (OPG) [Brojatsch et al., Cell, 87:845-855 (1996); Montgomery et al., Cell, 87:427-436 (1996); Marsters et al., J. Biol. Chem., 272:14029-14032 (1997); Simonet et al., Cell, 89:309-319 (1997)]. Unlike other known TNFR-like molecules, Simonet et al., supra, report that OPG contains no hydrophobic transmembrane-spanning sequence.

In Marsters et al., Curr. Biol., 6:750 (1996), investigators describe a full length native sequence human polypeptide, called Apo-3, which exhibits similarity to the TNFR family in its extracellular cysteine-rich repeats and resembles
5 TNFR1 and CD95 in that it contains a cytoplasmic death domain sequence [see also Marsters et al., Curr. Biol., 6:1669 (1996)]. Apo-3 has also been referred to by other investigators as DR3, wsl-1 and TRAMP. [Chinnaiyan et al., Science, 274:990 (1996); Kitson et al., Nature, 384:372 (1996); Bodmer et al., Immunity, 6:79 (1997)].

10 Pan et al. have disclosed another TNF receptor family member referred to as "DR4" [Pan et al., Science, 276:111-113 (1997)]. The DR4 was reported to contain a cytoplasmic death domain capable of engaging the cell suicide apparatus. Pan et al. disclose that DR4 is believed to be a receptor for the ligand known as Apo-2
15 ligand or TRAIL.

In Sheridan et al., Science, 277:818-821 (1997) and Pan et al., Science, 277:815-818 (1997), another molecule believed to be a receptor for the Apo-2 ligand (TRAIL) is described. That molecule is referred to as DR5 (it has also been alternatively referred to as
20 Apo-2). Like DR4, DR5 is reported to contain a cytoplasmic death domain and be capable of signaling apoptosis.

In Sheridan et al., supra, a receptor called DcR1 (or alternatively, Apo-2DcR) is disclosed as being a potential decoy receptor for Apo-2 ligand (TRAIL). Sheridan et al. report that DcR1
25 can inhibit Apo-2 ligand function *in vitro*. See also, Pan et al., supra, for disclosure on the decoy receptor referred to as TRID.

For a review of the TNF family of cytokines and their receptors, see Gruss and Dower, supra.

As presently understood, the cell death program contains
30 at least three important elements - activators, inhibitors, and effectors; in *C. elegans*, these elements are encoded respectively by three genes, *Ced-4*, *Ced-9* and *Ced-3* [Steller, Science, 267:1445 (1995); Chinnaiyan et al., Science, 275:1122-1126 (1997); Wang et al., Cell, 90:1-20 (1997)]. Two of the TNFR family members, TNFR1
35 and Fas/Apo1 (CD95), can activate apoptotic cell death [Chinnaiyan and Dixit, Current Biology, 6:555-562 (1996); Fraser and Evan, Cell;

85:781-784 (1996)]. TNFR1 is also known to mediate activation of the transcription factor, NF- κ B [Tartaglia et al., Cell, 74:845-853 (1993); Hsu et al., Cell, 84:299-308 (1996)]. In addition to some ECD homology, these two receptors share homology in their intracellular domain (ICD) in an oligomerization interface known as the death domain [Tartaglia et al., supra; Nagata, Cell, 88:355 (1997)]. Death domains are also found in several metazoan proteins that regulate apoptosis; namely, the Drosophila protein; Reaper, and the mammalian proteins referred to as FADD/MORT1, TRADD, and RIP [Cleaveland and Ihle, Cell, 81:479-482 (1995)]. Upon ligand binding and receptor clustering, TNFR1 and CD95 are believed to recruit FADD into a death-inducing signalling complex. CD95 purportedly binds FADD directly, while TNFR1 binds FADD indirectly via TRADD [Chinnaiyan et al., Cell, 81:505-512 (1995); Boldin et al., J. Biol. Chem., 270:387-391 (1995); Hsu et al., supra; Chinnaiyan et al., J. Biol. Chem., 271:4961-4965 (1996)]. It has been reported that FADD serves as an adaptor protein which recruits the Ced-3-related protease, MACH α /FLICE (caspase 8), into the death signalling complex [Boldin et al., Cell, 85:803-815 (1996); Muzio et al., Cell, 85:817-827 (1996)]. MACH α /FLICE appears to be the trigger that sets off a cascade of apoptotic proteases, including the interleukin-1 β converting enzyme (ICE) and CPP32/Yama, which may execute some critical aspects of the cell death programme [Fraser and Evan, supra].

It was recently disclosed that programmed cell death involves the activity of members of a family of cysteine proteases related to the *C. elegans* cell death gene, ced-3, and to the mammalian IL-1-converting enzyme, ICE. The activity of the ICE and CPP32/Yama proteases can be inhibited by the product of the cowpox virus gene, crmA [Ray et al., Cell, 69:597-604 (1992); Tewari et al., Cell, 81:801-809 (1995)]. Recent studies show that CrmA can inhibit TNFR1- and CD95-induced cell death [Enari et al., Nature, 375:78-81 (1995); Tewari et al., J. Biol. Chem., 270:3255-3260 (1995)].

As reviewed recently by Tewari et al., TNFR1, TNFR2 and CD40 modulate the expression of proinflammatory and costimulatory

cytokines, cytokine receptors, and cell adhesion molecules through activation of the transcription factor, NF- κ B [Tewari et al., Curr. Op. Genet. Develop., 6:39-44 (1996)]. NF- κ B is the prototype of a family of dimeric transcription factors whose subunits contain conserved Rel regions [Verma et al., Genes Develop., 9:2723-2735 (1996); Baldwin, Ann. Rev. Immunol., 14:649-681 (1996)]. In its latent form, NF- κ B is complexed with members of the I κ B inhibitor family; upon inactivation of the I κ B in response to certain stimuli, released NF- κ B translocates to the nucleus where it binds to specific DNA sequences and activates gene transcription. NF- κ B is induced by a variety of proinflammatory signals and cytokines including IL-1 and LPS acting through the Toll-like receptor TLR2 [Baeuerle et al., Ann. Rev. Immunol., 12:141-79 (1994); Verma et al., supra].

SUMMARY OF THE INVENTION

Applicants have identified a cDNA clone that encodes a novel polypeptide, designated in the present application as "DNA19355."

In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding DNA19355 polypeptide.

Optionally, the isolated nucleic acid comprises DNA encoding DNA19355 polypeptide having amino acid residues 1 to 177 or 52 to 177 of Fig. 1 (SEQ ID NO:1), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. The isolated nucleic acid may comprise the DNA19355 cDNA insert of the vector deposited as ATCC 209466, and particular the insert which includes the DNA sequence encoding DNA19355 polypeptide.

In another embodiment, the invention provides a vector comprising DNA encoding DNA19355 polypeptide. A host cell comprising such a vector is also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing DNA19355 polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of DNA19355 and recovering DNA19355 from the cell culture.

In another embodiment, the invention provides isolated DNA19355 polypeptide. In particular, the invention provides isolated native sequence DNA19355 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 177 or 52 to 177 of Figure 1 (SEQ ID NO:1). Optionally, the DNA19355 polypeptide is obtained or obtainable by expressing the polypeptide encoded by the cDNA insert of the vector deposited as ATCC 209466.

In another embodiment, the invention provides isolated DNA19355 polypeptide variants. The variants comprise polypeptides which have at least about 80% amino acid sequence identity with the deduced amino acid sequence of Fig. 1 (SEQ ID NO:1) or domain sequences identified herein, and preferably have activity(s) of native sequence or naturally-occurring DNA19355 polypeptide.

In another embodiment, the invention provides chimeric molecules comprising DNA19355 polypeptide fused to a heterologous polypeptide or amino acid sequence. An example of such a chimeric molecule comprises a DNA19355 fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In another embodiment, the invention provides an antibody which specifically binds to DNA19355 polypeptide. Optionally, the antibody is a monoclonal antibody.

In a still further embodiment, the invention provides diagnostic and therapeutic methods using DNA19355. For example, methods of inducing apoptosis in mammalian cancer cells are provided.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence (SEQ ID NO:2) of a cDNA for human DNA19355 and its derived amino acid sequence (SEQ ID NO:1).

Figure 2 shows an alignment and comparison of extracellular amino acid sequence of DNA19355 polypeptide with human Apo-2L, Fas/Apo1 ligand (CD95L), TNF-alpha and LT- α ; the respective

amino acid identities (%) are approximately 19.8, 19.0, 20.6, and 17.5.

Figure 3 shows a Northern blot analysis of DNA19355 mRNA expression in human tissues (identified adult and fetal tissues) and tumor cell lines (HL60 promyelocytic leukemia, HeLa S3 cervical carcinoma, K562 chronic myelogenous leukemia, MOLT4 lymphoblastic leukemia, Raji Burkitt's lymphoma, SW480 colorectal adenocarcinoma, A549 lung carcinoma, and G361 melanoma).

Figure 4 shows an analysis of soluble DNA19355 polypeptide by SDS-PAGE.

Figure 5 shows: (A) fluorescence images of Hoechst-stained nuclei from cells transfected with pRK5 (a); pRK5 encoding DNA19355 (b); pRK5 encoding Apo-2 ligand (c); pRK5 encoding DNA19355 plus pRK5 encoding CrmA (d); pRK5 encoding DNA19355 plus pRK5 encoding FADD-DN (e); pRK5 encoding Apo-2 ligand plus pRK5 encoding FADD-DN (f). (B) induction of apoptosis by transfected DNA19355 or Apo-2 ligand and effect of caspase inhibitors and FADD-DN.

Figure 6 shows the effect of DNA19355 on NF- κ B activity. Electrophoretic mobility shift analysis of NF- κ B activity in cells transfected with pRK5, or pRK5 encoding DNA19355, or pRK5 encoding Apo-2 ligand. In each case, the cells were co-transfected with pRK5 (left 3 lanes) or pRK5 encoding dominant-negative NIK (NIK-DN, right 3 lanes).

Figure 7 shows an alignment and comparison of the amino acid sequences for human GITR (hGITR) and murine GITR (mGITR). The three cysteine rich domains (CRD1, CRD2, and CRD3) and transmembrane region (TM) are shown.

Figure 8 shows the results of a co-precipitation assay described in Example 12 below. The autoradiograph of the SDS-PAGE gel revealed the hGITR-IgG molecule bound to the radioiodinated

DNA19355 polypeptide. Binding was not observed for the other immunoadhesin constructs identified.

Figure 9A shows the results of FACS analysis of transfected 293 cells assayed for binding to the identified receptors or ligand immunoadhesin constructs.

Figure 9B shows the results of FACS analysis of HUVEC cells assayed for binding to the identified receptor immunoadhesin constructs.

Figure 10 shows the results of a luciferase activity assay conducted to demonstrate NF- κ B activation by DNA19355/hGTR.

Figure 11 shows the results of an ELISA conducted to determine TNF-alpha levels in culture supernatants from primary T cells and monocytes/macrophages incubated with DNA19355 polypeptide.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. Definitions

For the purposes of this specification it will be clearly understood that the word "comprising" means "including but not limited to", and that the word "comprises" has a corresponding meaning.

The terms "DNA19355 polypeptide" and "DNA19355" when used herein encompass native sequence DNA19355 and DNA19355 variants (which are further defined herein). The DNA19355 may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The terms "DNA19355 polypeptide" and "DNA19355" when used herein refer to the same polypeptides referred to in the literature as "GLITTER". A "native sequence DNA19355" comprises a polypeptide having the same amino acid sequence as a DNA19355 derived from nature. Such native sequence DNA19355 can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence DNA19355" specifically encompasses naturally-occurring truncated, soluble or secreted forms of the DNA19355 (e.g., an extracellular domain sequence or soluble form),



naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the DNA19355. In one embodiment of the invention, the native sequence DNA19355 is a mature or full-length native sequence DNA19355 polypeptide comprising amino acids 1 to 177 of Fig. 1 (SEQ ID NO:1). Alternatively, the DNA19355 polypeptide comprises amino acids 52 to 177 of Fig. 1 (SEQ ID NO:1). Optionally, the DNA19355 polypeptide is obtained or obtainable by expressing the polypeptide encoded by the cDNA insert of the vector deposited as ATCC 209466.

The "DNA19355 extracellular domain" or "DNA19355 ECD" refers to a form of DNA19355 which is essentially free of the transmembrane and cytoplasmic domains of DNA19355. Ordinarily, DNA19355 ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. Optionally, DNA19355 ECD will comprise amino acid residues X to 177 of Fig. 1 (SEQ ID NO:1), wherein X is any one of amino acid residues 48 to 57 of Fig. 1 (SEQ ID NO:1). It will be understood by the skilled artisan that the transmembrane domain identified for the DNA19355 polypeptide of the present invention is identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain specifically mentioned herein.

"DNA19355 variant" means a DNA19355 as defined below having at least about 80% amino acid sequence identity with the DNA19355 having the deduced amino acid sequence shown in Fig. 1 (SEQ ID NO:1) for a full-length native sequence DNA19355 or the various domain sequences identified herein. Such DNA19355 variants include, for instance, DNA19355 polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the sequence of Fig. 1 (SEQ ID NO:1). DNA19355 variants include ECD fragments which include sequences having less than amino acid residues 52 to 177 of Fig. 1 (SEQ ID NO:1). Ordinarily, a DNA19355 variant will have at least about 80% or 85% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, and even more preferably at least about 95% amino acid

sequence identity with the amino acid sequence of Fig. 1 (SEQ ID NO:1).

"Percent (%) amino acid sequence identity" with respect to the DNA19355 sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the DNA19355 sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity.

Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

"Percent (%) nucleic acid sequence identity" with respect to the DNA19355 sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the DNA19355 sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising DNA19355, or a domain sequence thereof, fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, or which can be identified by some other agent, yet is short enough such that it does not interfere with activity of the DNA19355. The tag polypeptide preferably also is fairly unique so

that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 to about 50 amino acid residues (preferably, between about 10 to about 20 residues).

5 "Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural
10 environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup
15 sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the DNA19355 natural environment will not be present. Ordinarily, however,
20 isolated polypeptide will be prepared by at least one purification step.

An "isolated" DNA19355 nucleic acid molecule is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily
25 associated in the natural source of the DNA19355 nucleic acid. An isolated DNA19355 nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated DNA19355 nucleic acid molecules therefore are distinguished from the DNA19355 nucleic acid molecule as it exists in natural cells. However, an isolated
30 DNA19355 nucleic acid molecule includes DNA19355 nucleic acid molecules contained in cells that ordinarily express DNA19355 where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences
35 necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are

suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

5 Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably
10 linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a
15 secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

20 The term "antibody" is used in the broadest sense and specifically covers single anti-DNA19355 monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies) and anti-DNA19355 antibody compositions with polypeptidic specificity. The term "monoclonal antibody" as used herein refers to an antibody
25 obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

"Biologically active" and "desired biological activity"
30 for the purposes herein mean (1) having the ability to modulate apoptosis (either in an agonistic or stimulating manner or in an antagonistic or blocking manner) in at least one type of mammalian cell in vivo or ex vivo or (2) having the ability to induce or stimulate a proinflammatory response in at least one type of
35 mammalian cell in vivo or ex vivo.

The terms "apoptosis" and "apoptotic activity" are used in a broad sense and refer to the orderly or controlled form of cell death in mammals that is typically accompanied by one or more characteristic cell changes, including condensation of cytoplasm, loss of plasma membrane microvilli, segmentation of the nucleus, degradation of chromosomal DNA or loss of mitochondrial function. This activity can be determined and measured, for instance, by cell viability assays, FACS analysis or DNA electrophoresis, all of which are known in the art.

The terms "treating," "treatment," and "therapy" as used herein refer to curative therapy, prophylactic therapy, and preventative therapy.

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small-cell lung cancer, blastoma, gastrointestinal cancer, renal cancer, pancreatic cancer, glioblastoma, neuroblastoma, cervical cancer, ovarian cancer, liver cancer, stomach cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial cancer, salivary gland cancer, kidney cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, and various types of head and neck cancer.

The term "mammal" as used herein refers to any mammal classified as a mammal, including humans, cows, horses, dogs and cats. In a preferred embodiment of the invention, the mammal is a human.

II. Compositions and Methods of the Invention

A. DNA19355 Polypeptides

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as DNA19355. In particular, Applicants have identified and isolated cDNA encoding a DNA19355 polypeptide, as

disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that DNA19355 (shown in Figure 1 and SEQ ID NO:1) shares certain amino acid sequence identity with some members of the TNF family (see, e.g., Figure 2 and Example 1 below). As shown in the Examples below, DNA19355 polypeptide was found to have apoptotic activity and specific binding to G1TR. It was also found that DNA19355 stimulated secretion of TNF-alpha in primary T cells in vitro (Example 14) and infiltrate or influx of neutrophils in a guinea pig skin biopsy assay (such as described in Example 15), suggesting the role of DNA19355 in proinflammatory responses.

In addition to the full-length native sequence DNA19355 and soluble forms of DNA19355 described herein, it is contemplated that DNA19355 variants can be prepared. DNA19355 variants can be prepared by introducing appropriate nucleotide changes into the DNA19355 nucleotide sequence, or by synthesis of the desired DNA19355 polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the DNA19355, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence DNA19355 or in various domains of the DNA19355 described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the DNA19355 that results in a change in the amino acid sequence of the DNA19355 as compared with the native sequence DNA19355. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the DNA19355. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the DNA19355 with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino

acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity in any of the in vitro assays described in the Examples below.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the DNA19355 variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

B. Modifications of DNA19355

Covalent modifications of DNA19355 are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of the DNA19355 with an organic derivatizing agent that is capable of reacting with

selected side chains or the N- or C- terminal residues of the DNA19355. Derivatization with bifunctional agents is useful, for instance, for crosslinking DNA19355 to a water-insoluble support matrix or surface for use in the method for purifying anti-DNA19355 antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azido-salicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidyl-propionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the DNA19355 polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence DNA19355, and/or adding one or more glycosylation sites that are not present in the native sequence DNA19355.

Addition of glycosylation sites to the DNA19355 polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence DNA19355 (for O-linked glycosylation sites). The DNA19355 amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding

the DNA19355 polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the DNA19355 polypeptide is by chemical or enzymatic
5 coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the DNA19355
10 polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by
15 Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of DNA19355
20 comprises linking the DNA19355 polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

25 The DNA19355 of the present invention may also be modified in a way to form a chimeric molecule comprising DNA19355 fused to another, heterologous polypeptide or amino acid sequence. In one embodiment, such a chimeric molecule comprises a fusion of the DNA19355 with a tag polypeptide which provides an epitope to
30 which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the DNA19355. The presence of such epitope-tagged forms of the DNA19355 can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the DNA19355 to be
35 readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag.

In an alternative embodiment, the chimeric molecule may comprise a fusion of the DNA19355 with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule, such a fusion could be to the Fc region of an IgG molecule. In particular, the chimeric molecule may comprise a DNA19355 ECD fused to a His-tag molecule.

Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

The DNA19355 of the invention may also be modified in a way to form a chimeric molecule comprising DNA19355 fused to a leucine zipper. Various leucine zipper polypeptides have been described in the art. See, e.g., Landschulz et al., Science, 240:1759 (1988); WO 94/10308; Hoppe et al., FEBS Letters, 344:1991 (1994); Maniatis et al., Nature, 341:24 (1989). It is believed that use of a leucine zipper fused to DNA19355 may be desirable to assist in dimerizing or trimerizing soluble DNA19355 in solution. Those skilled in the art will appreciate that the leucine zipper may be fused at either the 5' or 3' end of the DNA19355 molecule.

C. Preparation of DNA19355

The description below relates primarily to production of DNA19355 by culturing cells transformed or transfected with a vector containing DNA19355 nucleic acid. It is, of course, contemplated

that alternative methods, which are well known in the art, may be employed to prepare DNA19355. For instance, the DNA19355 sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the DNA19355 may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length DNA19355.

1. Isolation of DNA Encoding DNA19355

DNA encoding DNA19355 may be obtained from a cDNA library prepared from tissue believed to possess the DNA19355 mRNA and to express it at a detectable level. Accordingly, human DNA19355 DNA can be conveniently obtained from a cDNA library prepared from human tissue. The DNA19355-encoding gene may also be obtained from a genomic library or by oligonucleotide synthesis.

Libraries can be screened with probes (such as antibodies to the DNA19355 or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding DNA19355 is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in

the art, and include the use of radiolabels like ³²P-labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

5 Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private
10 sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined through sequence alignment using computer software programs such as ALIGN, DNASTAR, and INHERIT which employ various algorithms to measure homology.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the
15 deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

20 2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for DNA19355 production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or
25 amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in
30 Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO₄ and electroporation. Depending on the host cell used, transformation is performed using standard
35 techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra,

or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635).

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for DNA19355-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism.

Suitable host cells for the expression of glycosylated DNA19355 are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line

transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol., 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

10

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding DNA19355 may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

25

The DNA19355 may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the DNA19355 DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter

30

35

described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may
5 be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more
10 selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning
15 vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or
20 tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to
25 take up the DNA19355 nucleic acid, such as DHFR or thymidine kinase.

An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene
30 present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12
35 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the DNA19355 nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., *Nature*, 275:615 (1978); Goeddel et al., *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (trp) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the tac promoter [deBoer et al., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding DNA19355.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

DNA19355 transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from

heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the DNA19355 by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the DNA19355 coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding DNA19355.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of DNA19355 in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled

probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence DNA19355 polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to DNA19355 DNA and encoding a specific antibody epitope.

5. Purification of Polypeptide

Forms of DNA19355 may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of DNA19355 can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify DNA19355 from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the DNA19355. Various methods of protein purification may be employed and such methods are

known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular DNA19355 produced.

D. Uses for DNA19355

Nucleotide sequences (or their complement) encoding DNA19355 have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. DNA19355 nucleic acid will also be useful for the preparation of DNA19355 polypeptides by the recombinant techniques described herein.

The full-length native sequence DNA19355 (Fig. 1; SEQ ID NO:2) gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate, for instance, still other genes (like those encoding naturally-occurring variants of DNA19355 or DNA19355 from other species) which have a desired sequence identity to the DNA19355 sequence disclosed in Fig. 1 (SEQ ID NO:2). Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequences including promoters, enhancer elements and introns of native sequence DNA19355. By way of example, a screening method will comprise isolating the coding region of the DNA19355 gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ³²P or ³⁵S, or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the DNA19355 gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

Nucleotide sequences encoding a DNA19355 can also be used to construct hybridization probes for mapping the gene which

encodes that DNA19355 and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as *in situ* hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

Screening assays can be designed to find lead compounds that mimic the biological activity of a native sequence DNA19355 or a ligand or receptor for DNA19355. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

Nucleic acids which encode DNA19355 or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding DNA19355 can be used to clone genomic DNA encoding DNA19355 in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding DNA19355. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for DNA19355 transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding DNA19355 introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased

expression of DNA encoding DNA19355. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression.

In accordance with this facet of the invention, an animal is
5 treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of DNA19355 can be
10 used to construct a DNA19355 "knock out" animal which has a defective or altered gene encoding DNA19355 as a result of homologous recombination between the endogenous gene encoding DNA19355 and altered genomic DNA encoding DNA19355 introduced into an embryonic cell of the animal. For example, cDNA encoding
15 DNA19355 can be used to clone genomic DNA encoding DNA19355 in accordance with established techniques. A portion of the genomic DNA encoding DNA19355 can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered
20 flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has
25 homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be
30 implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which
35 all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their

ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the DNA19355 polypeptide.

The DNA19355 polypeptides may also be employed in
5 diagnostic assays to, for instance, detect the presence of the receptor "GTR" in mammalian tissues. Such assays may be conducted using techniques known in the art or for example, using the binding assays described herein.

The DNA19355 polypeptides may further be employed as
10 immunogens to raise antibodies against DNA19355. Techniques and methods for generating antibodies are described below.

The DNA19355 polypeptides can also be employed therapeutically. For example, the DNA19355 polypeptides can be employed to induce apoptosis in mammalian cancer cells. Generally,
15 the methods for inducing apoptosis in mammalian cancer cells comprise exposing the cells to an effective (or apoptosis-inducing) amount of the DNA19355 polypeptide. Therapeutic application of DNA19355 polypeptide for the treatment of cancer is described in detail below.

In the methods for treating cancer, DNA19355 polypeptide
20 is administered to a mammal diagnosed as having cancer. It is of course contemplated that the DNA19355 polypeptide can be employed in combination with still other therapeutic compositions and techniques, including other apoptosis-inducing agents, chemotherapy,
25 radiation therapy and surgery.

The DNA19355 polypeptide is administered in an acceptable carrier, and preferably, a pharmaceutically-acceptable carrier. Suitable carriers and their formulations are described in Remington's Pharmaceutical Sciences, 16th Ed., 1980, Mack Publishing
30 Co., edited by Oslo et al. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the formulation to render the formulation isotonic. Examples of acceptable carriers include saline, Ringer's solution, and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and more
35 preferably, from about 7.4 to about 7.8. It will be apparent to those persons skilled in the art that certain carriers may be more

preferable depending upon, for instance, the route of administration and concentration of the DNA19355 polypeptide being administered.

The DNA19355 polypeptide may be administered to a mammal by injection (e.g., intravenous, intraperitoneal, subcutaneous, intramuscular), or by other methods such as infusion that ensure its delivery to the bloodstream in an effective form. It is also contemplated that the DNA19355 polypeptide can be administered by in vivo or ex vivo gene therapy.

Effective dosages and schedules for administering DNA19355 polypeptide may be determined empirically, and making such determinations is within the skill in the art. It is presently believed that an effective dosage or amount of DNA19355 polypeptide may range from about 1 microgram/kg to about 100 mg/kg of body weight or more per day. Interspecies scaling of dosages can be performed in a manner known in the art, e.g., as disclosed in Mordenti et al., *Pharmaceut. Res.*, 8:1351 (1991). Those skilled in the art will understand that the dosage of DNA19355 polypeptide that must be administered will vary depending on, for example, the mammal which will receive the DNA19355 polypeptide, the route of administration, and other drugs or therapies being administered to the mammal.

The one or more other therapies administered to the mammal may include but are not limited to, chemotherapy and/or radiation therapy, immunoadjuvants, cytokines, and antibody-based therapies. Examples include interleukins (e.g., IL-1, IL-2, IL-3, IL-6), leukemia inhibitory factor, interferons, erythropoietin, anti-VEGF antibody, and Her-2 antibody. Other agents known to induce apoptosis in mammalian cells may also be employed, and such agents include TNF-alpha, TNF-beta, CD30 ligand, 4-1BB ligand and Apo-1 ligand.

Chemotherapies contemplated by the invention include chemical substances or drugs which are known in the art and are commercially available, such as Doxorubicin, 5-Fluorouracil, etoposide, camptothecin, leucovorin, Cytosin arabinoside (Ara-C), Cyclophosphamide, Thiotepa, Busulfan, Cytosin, Taxol, methotrexate, Cisplatin, Melphalin, Vinblastine, and Carboplatin. Preparation and

dosing schedules for such chemotherapy may be used according to manufacturer's instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in Chemotherapy Service Ed., M.C.

5 Perry, Williams & Wilkins, Baltimore, MD (1992).

The chemotherapy is administered in an acceptable carrier, preferably a pharmaceutically-acceptable carrier, such as those described above. The mode of administration of the chemotherapy may be the same as employed for the DNA19355
10 polypeptide or it may be administered to the mammal via a different mode. For example, the DNA19355 polypeptide may be injected while the chemotherapy is administered orally to the mammal.

Radiation therapy can be administered to the mammal according to protocols commonly employed in the art and known to the
15 skilled artisan. Such therapy may include cesium, iridium, iodine or cobalt radiation. The radiation may be whole body irradiation, or may be directed locally to a specific site or tissue in or on the body. Typically, radiation therapy is administered in pulses over a period of time from about 1 to about 2 weeks. Optionally, the
20 radiation therapy may be administered as a single dose or as multiple, sequential doses.

The DNA19355 polypeptide and one or more other therapies may be administered to the mammal concurrently or sequentially. Following administration of DNA19355 polypeptide and one or more
25 other therapies to the mammal, the mammal's physiological condition can be monitored in various ways well known to the skilled practitioner. For instance, tumor mass may be observed physically, by biopsy, or by standard x-ray imaging techniques.

The modes and methods of administering DNA19355
30 polypeptide described above may also be used by the skilled practitioner to treat conditions whereby stimulation or induction of a proinflammatory response is desired.

E. Anti-DNA19355 Antibodies

The present invention further provides anti-DNA19355 antibodies. Exemplary antibodies include polyclonal, monoclonal,
5 humanized, bispecific, and heteroconjugate antibodies.

1. Polyclonal Antibodies

The DNA19355 antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to
10 the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the DNA19355 polypeptide or a fusion protein thereof.

15 It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed
20 include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

2. Monoclonal Antibodies

25 The DNA19355 antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or
30 other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

35 The immunizing agent will typically include the DNA19355 polypeptide or a fusion protein thereof. Generally, either

peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against DNA19355. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known

in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown in vivo as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison et al., supra] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

3. Humanized Antibodies

The DNA19355 antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab'), or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human

immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeven et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species.

In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985) and Boerner et al., J. Immunol., 147(1):86-95 (1991)).

4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the DNA19355, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, Nature, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., EMBO J., 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

5. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond.

Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimide and those disclosed, for example, in U.S. Patent No. 4,676,980.

5

F. Uses for DNA19355 Antibodies

The DNA19355 antibodies of the invention have various utilities. For example, DNA19355 antibodies may be used in diagnostic assays for DNA19355, e.g., detecting its expression in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, Monoclonal Antibodies: A Manual of Techniques, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , or ^{125}I , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., Nature, 144:945 (1962); David et al., Biochemistry, 13:1014 (1974); Pain et al., J. Immunol. Meth., 40:219 (1981); and Nygren, J. Histochem. and Cytochem., 30:407 (1982).

DNA19355 antibodies also are useful for the affinity purification of DNA19355 from recombinant cell culture or natural sources. In this process, the antibodies against DNA19355 are immobilized on a suitable support, such as a Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the DNA19355 to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the DNA19355, which is bound to the immobilized

antibody. Finally, the support is washed with another suitable solvent that will release the DNA19355 from the antibody.

5 The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their
10 entirety.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless
15 otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, Virginia.

20

EXAMPLE 1

Isolation of Human DNA19355

Methods described in Klein et al., PNAS, 93:7108-7113 (1996) were employed with the following modifications. Yeast transformation was performed with limiting amounts of transforming
25 DNA in order to reduce the number of multiple transformed yeast cells. Instead of plasmid isolation from the yeast followed by transformation of *E. coli* as described in Klein et al., supra, PCR analysis was performed on single yeast colonies. This was accomplished by restreaking the original sucrose positive colony
30 onto fresh sucrose medium to purify the positive clone. A single purified colony was then used for PCR using the following primers:
TGTAACGACGCGCCAGTTCTCTCAGAGAAACAAAGCAAAC (SEQ ID NO:7) and
CAGGAAACAGCTATGACCGAAGTGGACCAAAGGTCTATCGCTA (SEQ ID NO:8). The PCR primers are bipartite in order to amplify the insert and a small
35 portion of the invertase gene (allowing to determine that the insert

was in frame with invertase) and to add on universal sequencing primer sites.

A library of cDNA fragments derived from human HUVEC cells fused to invertase was transformed into yeast and transformants were selected on SC-URA media. URA and transformants were replica plated onto sucrose medium in order to identify clones secreting invertase. Positive clones were re-tested and PCR products were sequenced. The sequence of one clone, DNA1840, was determined to contain a signal peptide coding sequence.

Oligonucleotide primers and probes were designed using the nucleotide sequence of DNA1840. A full length plasmid library of cDNAs from human umbilical vein endothelial cells (HUVEC) was titrated and approximately 100,000 cfu were plated in 192 pools of 500 cfu/pool into 96-well round bottom plates. The pools were grown overnight at 37°C with shaking (200rpm). PCR was performed on the individual cultures using primers specific to DNA1840. Agarose gel electrophoresis was performed and positive wells were identified by visualization of a band of the expected size. Individual positive clones were obtained by colony lift followed by hybridization with ³²P-labeled oligonucleotide. These clones were characterized by PCR, restriction digest, and southern blot analyses.

A cDNA clone was sequenced in entirety. A nucleotide sequence of DNA19355 is shown in Figure 1 (SEQ ID NO:2). Clone DNA19355-1150 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 21-23 [Kozak et al., *supra*] (Fig. 1; SEQ ID NO:2). The predicted polypeptide precursor is 177 amino acids long and has a calculated molecular weight of approximately 20,308 daltons. Hydropathy analysis suggests a type II transmembrane protein typology, with a putative cytoplasmic region (amino acids 1-25); transmembrane region (amino acids 26-51); and extracellular region (amino acids 52-177). Two potential N-linked glycosylation sites have been identified at position 129 (Asn) and position 161 (Asn) of the sequence shown in Fig. 1 (SEQ ID NO:1). Clone DNA19355-1150 has been deposited with ATCC and is assigned ATCC deposit no. 209466. DNA19355 polypeptide is obtained or obtainable by expressing the molecule encoded by the

cDNA insert of the deposited ATCC 209466 vector. Digestion of the vector with XbaI and NotI restriction enzymes will yield a 1411 bp fragment and 668 bp fragment.

Based on a BLAST and FastA sequence alignment analysis (using the ALIGN computer program) of extracellular sequence, DNA19355 shows amino acid sequence identity to several members of the TNF cytokine family, and particularly, to human Apo-2L (19.8%), Fas/Apo1-ligand (19.0%), TNF-alpha (20.6%) and Lymphotoxin-alpha (17.5%) (see Figure 2). Most of the amino acid sequence identity is found in the regions corresponding to the beta-strands in the crystal structure of TNF-alpha [Banner et al., *Cell*, 73:431-435 (1993); Eck et al., *J. Biol. Chem.*, 264:17595-605 (1989); Lewit-Bentley et al., *J. Mol. Biol.*, 199:389-92 (1988)]. The sequence of strand C is especially conserved in all members of the family (see Figure 2). The sequence between the putative transmembrane domain and the first beta-strand of the DNA19335 polypeptide is relatively short, including 5 residues, as compared to about 30 to about 80 residues in TNF-alpha, CD95L or Apo-2 ligand.

EXAMPLE 2

Northern Blot Analysis

Expression of DNA19355 mRNA in human tissues and tumor cell lines was examined by Northern blot analysis (see Figure 3). Human RNA blots were hybridized to an approximately 700 bp-long ³²P-labeled DNA probe generated by digestion of the pRK5 plasmid encoding full-length DNA19355 cDNA with Xba-I; this probe corresponds to the entire coding sequence plus some flanking 5' and 3' sequences.

Human fetal, adult, or cancer cell line mRNA blots (Clontech) were incubated with the DNA probe in hybridization buffer (5X SSPE; 2X Denhardt's solution; 100 mg/mL denatured sheared salmon sperm DNA; 50% formamide; 2% SDS) for 60 hours at 42°C. The blots were washed several times in 2X SSC; 0.05% SDS for 1 hour at room temperature, followed by a 30 minute wash in 0.1X SSC; 0.1% SDS at 50°C. The blots were developed after overnight exposure by phosphorimager analysis (Fuji).

As shown in Figure 3, a predominant mRNA transcript of about 3.2 kB was detected in fetal kidney and lung, and in adult small intestine. Expression was also detected in 6 out of 8 human tumor cell lines tested, which showed about the same 3.2 kB transcript, as well as weaker expression of about 1.5 and about 5 kB transcripts.

The results indicate that the mRNA expression of the DNA19355 polypeptide is relatively restricted in normal tissues, but is markedly elevated in tumor cell lines from lymphoid as well as non-lymphoid origin.

EXAMPLE 3

Expression of DNA19355 in E. coli

The DNA sequence (of Fig. 1; SEQ ID NO:2) encoding an extracellular region of the DNA19355 polypeptide (amino acids 52 to 177 of Fig. 1; SEQ ID NO:1) was amplified with PCR primers containing flanking NdeI and XbaI restriction sites, respectively: forward: 5'-GAC GAC AAG CAT ATG TTA GAG ACT GCT AAG GAG CCC TG-3' (SEQ ID NO:3); reverse: 5'-TAG CAG CCG GAT CCT AGG AGA TGA ATT GGG GATT-3' (SEQ ID NO:4). The PCR was digested and cloned into the NdeI and XbaI sites of plasmid pET19B (Novagen) downstream and in frame of a Met Gly His₁₂ sequence followed by a 12 amino acid enterokinase cleavage site (derived from the plasmid): Met Gly His His His His His His His His His Ser Ser Gly His Ile Asp Asp Asp Asp Lys His Met (SEQ ID NO:5).

The resulting plasmid was used to transform E. coli strain JM109 (ATCC 53323) using the methods described in Sambrook et al., *supra*. Transformants were identified by PCR. Plasmid DNA was isolated and confirmed by restriction analysis and DNA sequencing.

Selected clones were grown overnight in liquid culture medium LB supplemented with antibiotics. The overnight culture was subsequently used to inoculate a larger scale culture. The cells were grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells were harvested by centrifugation. The cell pellet obtained by

the centrifugation was solubilized using a microfluidizer in a buffer containing 0.1M Tris, 0.2M NaCl, 50 mM EDTA, pH 8.0. The solubilized DNA19355 protein was purified using Nickel-sepharose affinity chromatography.

5 The DNA19355 protein was analyzed by SDS-PAGE followed by Western Blot with nickel-conjugated horseradish peroxidase followed by ECL detection (Boehringer Mannheim). Three predominant protein bands were detected, which corresponded in size to monomeric, homodimeric, and homotrimeric forms of the protein
10 (Figure 4). It is believed based on this result that in its native form, in the absence of SDS denaturation, the soluble DNA19355 protein is capable of forming homotrimers.

15

EXAMPLE 4

Apoptotic Activity of DNA19355

The PRK5 plasmid encoding the full-length DNA19355 protein, or empty PRK5 plasmid, or PRK5 encoding full-length human Apo-2 ligand (Apo-2L) was transiently transfected into human 293
20 cells (10^6 cells/10 cm dish) by calcium phosphate precipitation. In some cases, the cells were co-transfected with a PRK5 plasmid encoding the poxvirus-derived caspase inhibitor CrmA, or a dominant-negative mutant form of death adaptor protein FADD (FADD-DN), which mediates death signaling by Fas/Apo1 and TNFR1. Sixteen hours
25 later, the cells were stained with Hoechst 33342 dye (10 μ g/ml), and apoptotic or normal nuclei were counted under a Leica fluorescence microscope equipped with Hoffmann optics. In some cases, the caspase inhibitor z-VAD-fmk (Research Biochemicals) (200 μ M) was added to the dishes immediately after transfection.

30 As shown in Fig. 5, transfection by DNA19355 resulted in a substantial increase in the level of apoptosis as compared with PRK5, similar to the increase observed with Apo-2L, a well-established apoptosis inducer. The increase in apoptosis induced by DNA19355 was blocked by CrmA or by z-VAD-fmk, indicating the
35 involvement of caspases in this effect. In addition, the increase in apoptosis induced by DNA19355, but not by Apo-2L, was blocked by

FADD-DN, indicating that the FADD adaptor protein plays an essential role in transmitting the death signal from DNA19355 to the caspase machinery.

5

EXAMPLE 5

Activation of NF- κ B by DNA19355

The pRK5 plasmid encoding the full-length DNA19355 protein, or empty pRK5 plasmid, or pRK5 encoding full-length human
10 Apo-2L was transiently transfected into human 293 cells (10⁶ cells/10 cm dish) by calcium phosphate precipitation. The cells were co-transfected with empty pRK5 plasmid, or pRK5 plasmid encoding a dominant-negative, kinase deficient, mutant form of the serine/threonine kinase NIK (NIK-DN), which mediates NF- κ B
15 activation by TNF [Malinin et al., *Nature*, 385:540-544 (1997)]. Sixteen hours later, the cells were harvested, nuclear extracts were prepared, and 1 μ g of nuclear protein was reacted with a ³²P-labeled NF- κ B-specific synthetic oligonucleotide probe ATCAGGGACTTTCGCTGGGGACTTTCG (SEQ ID NO:6) [see, also, MacKay et
20 al., *J. Immunol.*, 153:5274-5284 (1994)].

As shown in Fig. 6, transfection by DNA19355 induced significant NF- κ B activation, as measured by an electrophoretic mobility shift assay [Marsters et al. *PNAS*, 92:5401-5405 (1995)]; the level of activation was greater than the level obtained with
25 Apo-2L. Co-transfection with NIK-DN substantially reduced NF- κ B activation by DNA19355, but not activation by Apo-2L. This result suggests that like TNF- α , DNA19355 activates NF- κ B through a signaling pathway that involves the NIK protein.

30

EXAMPLE 6

Expression of DNA19355 in mammalian cells

This example illustrates preparation of a form of DNA19355 by recombinant expression in mammalian cells.

35 The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the

DNA19355 DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the DNA19355 DNA using ligation methods such as described in Sambrook et al., supra. The resulting vector is called pRK5-DNA19355.

5 In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 µg pRK5-DNA19355 DNA is mixed with about 1 µg DNA encoding
10 the VA RNA gene [Thimmappaya et al., Cell, 31:543 (1982)] and dissolved in 500 µl of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 µl of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the
15 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

20 Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 µCi/ml ³⁵S-cysteine and 200 µCi/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15%
25 SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of DNA19355 polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

30 In an alternative technique, DNA19355 may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac et al., Proc. Natl. Acad. Sci., 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 µg pRK5-DNA19355 DNA is added. The cells are first
35 concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell

pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 µg/ml bovine insulin and 0.1 µg/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed DNA19355 can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, DNA19355 can be expressed in CHO cells. The pRK5-DNA19355 can be transfected into CHO cells using known reagents such as CaPO₄ or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ³⁵S-methionine. After determining the presence of DNA19355 polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed DNA19355 can then be concentrated and purified by any selected method.

Epitope-tagged DNA19355 may also be expressed in host CHO cells. The DNA19355 may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged DNA19355 insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged DNA19355 can then be concentrated and purified by any selected method, such as by Ni²⁺-chelate affinity chromatography.

EXAMPLE 7Expression of DNA19355 in Yeast

The following method describes recombinant expression of
5 DNA19355 in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of DNA19355 from the ADH2/GAPDH promoter, DNA encoding DNA19355, a selected signal peptide and the promoter is inserted into suitable restriction
10 enzyme sites in the selected plasmid to direct intracellular expression of DNA19355. For secretion, DNA encoding DNA19355 can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, the yeast alpha-factor secretory signal/leader sequence, and linker sequences (if needed) for expression of
15 DNA19355.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10%
20 trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant DNA19355 can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected
25 cartridge filters. The concentrate containing DNA19355 may further be purified using selected column chromatography resins.

EXAMPLE 8Expression of DNA19355 in Baculovirus-infected Insect Cells

The following method describes recombinant expression of
30 DNA19355 in insect cells.

The DNA19355 is fused upstream of an epitope tag contained with a baculovirus expression vector. Such epitope tags
35 include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids

derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the DNA19355 or the desired portion of the DNA19355 (such as a sequence encoding an extracellular domain) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression is performed as described by O'Reilley et al., *Baculovirus expression vectors: A laboratory Manual*, Oxford:Oxford University Press (1994).

Expressed poly-his tagged DNA19355 can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., *Nature*, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% Glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% Glycerol, pH 7.8) and filtered through a 0.45 µm filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% Glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed

by SDS-PAGE and silver staining or western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₆-tagged DNA19355 are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Pc tagged) DNA19355 can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

10

EXAMPLE 9

Preparation of Antibodies that Bind DNA19355

This example illustrates preparation of monoclonal antibodies which can specifically bind DNA19355.

Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified DNA19355, fusion proteins containing DNA19355, and cells expressing recombinant DNA19355 on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the DNA19355 immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect DNA19355 antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of DNA19355. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected

murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against DNA19355. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against DNA19355 is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-DNA19355 monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein-G can be employed.

EXAMPLE 10

Use of DNA19355 as a hybridization probe

The following method describes use of a nucleotide sequence encoding DNA19355 as a hybridization probe.

DNA comprising the coding sequence of DNA19355 (as shown in Figure 1, SEQ ID NO:2) is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of DNA19355) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled DNA19355-derived probe to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20

hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNA's having a desired sequence identity with the DNA encoding full-length native sequence DNA19355 can then be identified
5 using standard techniques known in the art.

EXAMPLE 11

Chromosomal Mapping

10 Chromosomal localization of the human DNA19355 gene was examined by radiation hybrid (RH) panel analysis. RH mapping was performed by PCR using a mouse-human cell radiation hybrid panel (Research Genetics) and primers based on the coding region of the DNA19355 cDNA [Gelb et al., Hum. Genet., 98:141 (1996)]. Analysis
15 of the PCR data using the Stanford Human Genome Center Database indicated that DNA19355 is linked to the STS marker D1S2790 and to Genethon marker AFMb352xe9, and maps to the human chromosome 1q23. Notably, CD95L also maps to chromosome 1q23 [Takahashi et al., Int. Immunol., 6:1567-1574 (1994); whereas OX40 ligand maps to chromosome
20 1q25 [Baum et al., EMBO J., 13:3992-4001 (1994)]. Accordingly, these TNF family members may have arisen by duplication and divergence of a common ancestral gene.

EXAMPLE 12

25 Binding Specificity of DNA19355 Polypeptide for Human GITR Receptor

Assays were conducted to determine whether the DNA19355 polypeptide interacts and specifically binds with a human homolog of the receptor molecule referred to as 'GITR'. A murine GITR (mGITR) polypeptide was described in Nocentini et al., Proc. Natl. Acad. Sci., 94:6216-6221 (1997). What are believed to be human
30 homologs of the mGITR have been described. An amino acid sequence for a full length human GITR (hGITR) is shown in SEQ ID NO:4 in PCT WO 98/06842, published February 19, 1998. A comparison of the hGITR and mGITR amino acid sequences is shown in Figure 7.

35 To test for binding, a soluble immunoglobulin fusion protein (immunoadhesin) which included the hGITR extracellular

domain (see amino acids 1-167 of Figure 7) was expressed in insect cells. The hGITR ECD was expressed as a C-terminus IgG-Fc tagged form in insect cells using Baculovirus (as described in Example 8 above).

5 A soluble DNA19355 polypeptide was also prepared by expressing the ECD in *E. coli* cells (as described in Example 3 above). The soluble DNA19355 ECD molecule was then labeled with ¹²⁵I. For comparison, immunoadhesin constructs were also made of the following TNF receptor family members: CD95, DR4, DR5, TNFR1, TNFR2, 10 and Apo-3. CD95, DR4, DR5, TNFR1, TNFR2, and Apo-3 immunoadhesins were prepared by fusing each receptor's ECD to the hinge and Fc portion of human IgG, as described previously for TNFR1 [Ashkenazi et al., *Proc. Natl. Acad. Sci.*, 88:10535-10539 (1991)]. The respective TNF receptor family members are described (and relevant 15 references cited) in the Background of the Invention section.

For the co-precipitation assay, each immunoadhesin (5 microgram) was incubated with ¹²⁵I-labeled soluble DNA19355 polypeptide (1 microgram) for 1 hour at 24°C, followed by protein A-sepharose for 30 minutes on ice. The reaction mixtures were spun 20 down and washed several times in PBS, boiled in SDS-PAGE buffer containing 20 mM dithiothreitol and then resolved by SDS-PAGE and autoradiography.

The results are shown in Figure 8. The position of the molecular weight markers (kDa) are indicated in the figure. The 25 hGITR-IgG bound to the radioiodinated soluble DNA19355 polypeptide. However, the hGITR-IgG did not bind to the immunoadhesin constructs of CD95, DR4, DR5, TNFR1, TNFR2, or Apo-3.

In another assay, human 293 cells were transiently transfected with DNA19355 and the ability of receptor immunoadhesin 30 constructs for hGITR, TNFR1, HVEM, and DcR1 to bind to those transfected cells was determined by FACS analysis. The 293 cells were maintained in high glucose DMEM media supplemented with 10% fetal bovine serum (FBS), 2mM glutamine, 100 microgram/ml penicillin, and 100 microgram/ml streptomycin. The transfected 35 cells (1×10^5) were incubated for 60 minutes at 4°C in 200 microliters 2% FBS/PBS with 1 microgram of the respective receptor

or ligand immunoadhesin. The cells were then washed with 2% FBS/PBS, stained with R-phycoerythrin-conjugated goat anti-human antibody (Jackson ImmunoResearch, West Grove, PA). Next, the cells were analyzed by FACS. To test the binding of the respective immunoadhesins to the transiently transfected cells, an expression vector (pRK5-CD4; Smith et al., *Science*, 328:1704-1707 (1987)) for CD4 was co-transfected with DNA19355 expression vector (see Example 3). FITC-conjugated anti-CD4 (Pharmingen, San Diego, CA) was then used to identify and gate the transfected cell population in the FACS analysis.

As shown in Figure 9A, the hGITR-IgG bound specifically to the surface of cells transfected with the expression plasmid encoding the full length DNA19355. No such binding was observed for the TNFR1, HVEM or DcR1. The hGITR-IgG did not bind to the cells transfected with a control plasmid (data not shown).

The results demonstrate a specific binding interaction of the DNA19355 polypeptide with hGITR and that the DNA19355 polypeptide does not interact with any of the other TNF receptor family members tested.

The DNA19355 polypeptide was identified in a human umbilical vein endothelial cell (HUVEC) library, and the DNA19355 polypeptide transcripts are readily detectable in HUVEC by RT-PCR (data not shown). A FACS analysis assay was conducted to examine whether specific binding of hGITR-IgG could be demonstrated with HUVEC by FACS analysis. HUVEC cells were purchased from Cell Systems (Kirkland, WA) and grown in a 50:50 mix of Ham's F12 and Low Glucose DMEM media containing 10% fetal bovine serum, 2 mM L-glutamine, 10 mM Hepes, and 10 ng/ml basic FGF. Cells were FACS sorted with PBS, hGITR-IgG, TNFR1-IgG or Fas-IgG as a primary antibody and goat anti-human F(ab')₂ conjugated to phycoerythrin (CalTag, Burlingame, CA).

It was found that hGITR-IgG specifically bound to HUVEC. (See Figure 9B). Neither the Fas-IgG nor the TNFR1-IgG exhibited specific binding to the HUVEC cells.

EXAMPLE 13Activation of NF- κ B by DNA19355

An assay was conducted to determine whether DNA19355/hGITR induces NF- κ B activation by analyzing expression of a reporter gene driven by a promoter containing a NF- κ B responsive element from the E-selectin gene.

Human 293 cells (2×10^5) were transiently transfected by calcium phosphate transfection with 0.5 microgram of the firefly luciferase reporter plasmid pGL3.BLAK.tk [Yang et al., Nature, 395:284-288 (1998)] and 0.05 microgram of the Renilla luciferase reporter plasmid (as internal transfection control) (Pharmacia), as well as the indicated additional expression vectors for DNA19355 and hGITR (described above) (0.1 microgram hGITR; 0.5 microgram for other expression vectors), and carrier plasmid pRK50 to maintain constant DNA between transfections. After 24 hours, the cells were harvested and luciferase activity was assayed as recommended by the manufacturer (Pharmacia). Activities were normalized for differences in transfection efficiency by dividing firefly luciferase activity by that of Renilla luciferase and were expressed as activity relative to that seen in the absence of added expression vectors.

As shown in Figure 10, overexpression of hGITR resulted in significant gene activation, and the observed result was enhanced by co-expression of both DNA19355 and hGITR.

EXAMPLE 14Stimulation of TNF-alpha Production

An assay was conducted to examine production of TNF-alpha and IL-1beta from isolated primary T cells and macrophages in response to stimulation by DNA19355 polypeptide.

Primary T cells or monocyte/macrophages were isolated from human donors. The primary human T cells were isolated from whole blood by a T cell enrichment column (R & D Systems). The monocytes/macrophages were isolated from whole blood by adherence to a tissue culture flask. The respective isolated cells were then

treated for 24 hours with the DNA19355 immunoadhesin (see Example 3 above) at 5 microgram/ml in RPMI 1640 medium containing 10% FBS. TNF-alpha levels in the culture supernatants were then determined by ELISA (R & D Systems; according to manufacturer's instructions).

5 The results are illustrated in Figure 11. The DNA19355 polypeptide induced about a 20-fold increase in secreted TNF-alpha levels from the T cells, but did not affect TNF-alpha release or IL-1beta release from macrophages (data not shown). The induced TNF-alpha production from the human T cells suggests that DNA19355
10 polypeptide/hGTR contribute to a proinflammatory response.

EXAMPLE 15

15 Guinea Pig Skin Biopsy Assay

An *in vivo* assay is conducted to determine the activity of a candidate molecule in proinflammatory responses. Specifically, a candidate molecule (such as DNA19355 polypeptide) is injected in guinea pigs and skin biopsies from the treated animal are analyzed
20 for polymorphonuclear/mononuclear cell infiltrate or eosinophil infiltrate.

The guinea pigs are anesthetized with Ketamine (75-80 mg/kg plus 5 mg/kg Xylazine) intramuscularly. The candidate molecule is then injected into skin on the back of the animal at 16
25 sites (100 microliter per site intradermally). Approximately 1 ml of Evans blue dye/PBS is injected intracordially.

Blemishes at the injection sites are measured (mm diameter) at 1 hour and 6 hours. The guinea pigs are sacrificed at 6 hours after skin injection. Skin samples at the injection sites
30 are excised and fixed in paraformaldehyde. The tissues are then prepared for histological evaluation using standard staining techniques. Analysis of the tissues includes characterizing cell type in the inflammatory infiltrate and evaluating the perivascular infiltrate.

35

Deposit of Biological Material

The following materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, Virginia USA (ATCC):

<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
DNA19355-1150	209466	November 18, 1997

10 This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of
15 deposit. The deposit will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying
20 open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular
25 reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same.
30 Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

The foregoing written specification is considered to be
35 sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended

as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is
5 inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to
10 those skilled in the art from the foregoing description and fall within the scope of the appended claims.

Applicant's or agent's file reference number	11669.26W002	International application No.	
--	--------------	-------------------------------	--

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page _____ line _____	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, Virginia USA	
Date of deposit November 18, 1997	Accession Number
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau (specify the general nature of the indications e.g. "Accession Number of Deposit")	
For receiving Office use only	
<input checked="" type="checkbox"/> This sheet was received with the international application I 11669.26W002 Authorized officer: [Signature] (799) 265-5747	
For International Bureau use only	
<input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer	

EDITORIAL NOTE 14180/99

**SEQUENCE LISTING PAGES 1 TO 4 ARE PART OF THE
DESCRIPTION AND ARE FOLLOWED BY CLAIM PAGES 63 TO 68.**

SEQUENCE LISTING

<110> GENENTECH INC.

<120> DNA19355 POLYPEPTIDE, A TUMOR NECROSIS FACTOR HOMOLOG

<130> 11669.26W001

<140> NEW FILING

<141> 1998-11-18

<150> 60/069,661

<151> 1997-12-12

<150> 60/065,635

<151> 1997-11-18

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 177

<212> PRT

<213> Homo sapiens

<400> 1

Met	Cys	Leu	Ser	His	Leu	Glu	Asn	Met	Pro	Leu	Ser	His	Ser	Arg	Thr
1				5					10					15	

Gln	Gly	Ala	Gln	Arg	Ser	Ser	Trp	Lys	Leu	Trp	Leu	Phe	Cys	Ser	Ile
			20					25					30		

Val	Met	Leu	Leu	Phe	Leu	Cys	Ser	Phe	Ser	Trp	Leu	Ile	Phe	Ile	Phe
	35						40					45			

Leu	Gln	Leu	Glu	Thr	Ala	Lys	Glu	Pro	Cys	Met	Ala	Lys	Phe	Gly	Pro
	50					55					60				

Leu	Pro	Ser	Lys	Trp	Gln	Met	Ala	Ser	Ser	Glu	Pro	Pro	Cys	Val	Asn
	65				70					75				80	

Lys	Val	Ser	Asp	Trp	Lys	Leu	Glu	Ile	Leu	Gln	Asn	Gly	Leu	Tyr	Leu
			85					90					95		

Ile	Tyr	Gly	Gln	Val	Ala	Pro	Asn	Ala	Asn	Tyr	Asn	Asp	Val	Ala	Pro
			100					105					110		

Phe Glu Val Arg Leu Tyr Lys Asn Lys Asp Met Ile Gln Thr Leu Thr
115 120 125

Asn Lys Ser Lys Ile Gln Asn Val Gly Gly Thr Tyr Glu Leu His Val
130 135 140

Gly Asp Thr Ile Asp Leu Ile Phe Asn Ser Glu His Gln Val Leu Lys
145 150 155 160

Asn Asn Thr Tyr Trp Gly Ile Ile Leu Leu Ala Asn Pro Gln Phe Ile
165 170 175

Ser

<210> 2

<211> 1964

<212> DNA

<213> Homo sapiens

<400> 2

cagctctcat ttctccaaaa atgtgtttga gccacttgga aaatatgcct ttaagccatt 60
caagaactca aggagctcag agatcatcct ggaagctgtg gctcttttgc tcaatagtta 120
tggtgtatt tctttgctcc ttcagttggc taatctttat tttctccaa ttagagactg 180
ctaaggagcc ctgtatggct aagtttgac cattaccctc aaaatggcaa atggcatctt 240
ctgaacctcc ttgcgtgaat aaggtgtctg actggaagct ggagatactt cagaatggct 300
tatatttaatt ttatggccaa gtggctccca atgcaaaacta caatgatgta gctccttttg 360
aggtgcggct gtataaaaaa aaagacatga taaaaactct aacaaacaaa tctaaaatcc 420
aaaatgtagg agggacttat gaattgcatg ttggggacac catagacttg atattcaact 480
ctgagcatca ggttctaaaa aataatacat actggggtat cattttacta gcaaaccctc 540
aattcatctc ctagagactt gatttgatct cctcattccc ttcagcacat gttaggtgct 600
cagtggttgg attggagggg gaagatattc aatttctaga gtttgtctgt ctacaaaaat 660
caacacaaac agaactcctc tgcacgtgaa ttttcatcta tcatgcctat ctgaaagaga 720
ctcaggggaa gagccaaaga cttttggttg gatctgcaga aatacttcat taatccatga 780
taaaacaaat atggatgaca gaggacatgt gcttttcaaa gaatctttat ctaattcttg 840
aattcatgag tggaaaaatg gagttctatt cccatggaag atttacctgg tatgcaaaaa 900
ggatctgggg cagtgcctg gctttgttct catattcttg ggctgctgta attcattctt 960
ctcactactcc catcttctga gacctccca ataaaaagta gactgatagg atggccacag 1020
atatgcctac cataccctac tttagatattg gtggtgttag aagataaaga acaatctgag 1080
aactattgga atagaggtac aagtggcata aaatggaatg tacgctatct ggaatttctt 1140
cttggtttta tcttctcag gatgcagggt gctttaaaaa gccttatcaa aggagtcatt 1200
ccgaacctc acgtagagct ttgtgagacc ttactgttg tgtgtgtgtc taaacattgc 1260
taattgtaaa gaaagagtaa ccattagtaa tcattaggtt taacccaga atggtattat 1320
cattactgga ttatgtcatg taatgattta gtatttttag ctactttcc acagtttgca 1380
aagtgtcttc gtaaaacagt tagcaattct atgaagttaa ttgggcaggc atttggggga 1440
aaattttagt gatgagaatg tgatagcata gcatagccaa ctttctcaa ctcataggac 1500
aagtactac aagaggcaat gggtagtccc ctgcattgca ctgtctcagc tttagaattg 1560

ttattttctgc tategtgta taagactcta aaacttagcg aattcacttt tcaggaagca 1620
 tattccccctt tagcccaagg tgagcagagt gaagctacaa cagatctttc cttaccagc 1680
 acactttttt ttttttttcc tgctgaatc agggagatcc aggatgtgt tcaggccaaa 1740
 tcccaaccaa attccccctt tcaactttgca gggcccatct tagtcaaatg tgctaacttc 1800
 taaaataata aatagcacta attcaaaatt ttggaatct taaattagct acttgcnngt 1860
 tgcttggtga aaggnatata atgattacat tgtaaacaaa tttaaaatat ttatggatat 1920
 ttgtgaaaag ctgcattatg ttaaataata ttacatgtaa agct 1964

<210> 3
 <211> 38
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism:Unknown

<400> 3
 gacgacaagc atatgttaga gactgctaag gagccctg 38

<210> 4
 <211> 34
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism:Unknown

<400> 4
 tagcagccgg atcctaggag atgaattggg gatt 34

<210> 5
 <211> 24
 <212> PRT
 <213> Unknown

<220>
 <223> Description of Unknown Organism:Unknown

<400> 5
 Met Gly His His His His His His His His His His Ser Ser Gly His
 1 5 10 15

Ile Asp Asp Asp Asp Lys His Met
 20

<210> 6
 <211> 29

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism:Unknown

<400> 6

atcagggact ttccgctggg gactttccg

29

<210> 7

<211> 42

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism:Unknown

<400> 7

tgtaaaacga cggccagttt ctctcagaga aacaagcaaa ac

42

<210> 8

<211> 43

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism:Unknown

<400> 8

caggaaacag ctatgaccga agtggaccaa aggtctatcg cta

43

- 63 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An isolated nucleic acid molecule comprising DNA encoding a DNA19355 polypeptide consisting of amino acid residues X to 177 of Fig. 1 (SEQ ID NO:1), wherein X is any one of amino acid residues 48 to 57 of Fig. 1 (SEQ ID NO:1).
2. The nucleic acid molecule of claim 1, comprising DNA encoding a DNA19355 polypeptide consisting of amino acid residues 52 to 177 of Fig. 1 (SEQ ID NO:1).
3. An isolated nucleic acid molecule comprising a DNA encoding a polypeptide having at least 80% amino acid sequence identity with native sequence DNA19355 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1), wherein said encoded polypeptide has the ability to
 - i) induce apoptosis in a mammalian cell,
 - ii) activate NF-KB in a mammalian cell,
 - iii) bind GITR receptor, or
 - iv) stimulate mammalian T cells to secrete TNF- α ; andwherein the polypeptide includes amino acid residues Y to 8 of SEQ ID NO:1, and wherein Y is any one of amino acid residues 1 to 8 of SEQ ID NO:1.
4. The nucleic acid molecule of claim 3, wherein said DNA encodes a polypeptide having at least 90% amino acid sequence identity with native sequence DNA19355 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1).
5. The nucleic acid molecule of claim 3, wherein said DNA encodes a polypeptide having at least 95% amino acid sequence identity with native sequence DNA19355 polypeptide consisting of amino acid residues 1 to 177 of



- 64 -

Fig. 1 (SEQ ID NO:1).

6. An isolated nucleic acid molecule comprising a DNA encoding a polypeptide having at least 96% amino acid sequence identity with native sequence DNA19355.
- 5 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1), wherein said encoded polypeptide has the ability to
- i) induce apoptosis in a mammalian cell,
 - 10 ii) activate NF- κ B in a mammalian cell,
 - iii) bind GITR receptor, or
 - iv) stimulate mammalian T cells to secrete TNF- α .
- 15 7. A vector comprising the nucleic acid of any one of claims 3 to 6.
8. The vector of claim 7, operably linked to control sequences recognized by a host cell transformed with the
- 20 vector.
9. A host cell comprising the vector of claim 7.
10. The host cell of claim 9, which is a CHO cell.
- 25 11. The host cell of claim 9, which is an *E. coli* cell.
12. The host cell of claim 9, which is a yeast cell.
- 30 13. A process for producing DNA19355 polypeptides, comprising culturing the host cell of any one of claims 9 to 12 under conditions suitable for expression of the polypeptide, and recovering the polypeptide from the cell
- 35 culture.
14. The isolated nucleic acid molecule of claim 3,



- 65 -

comprising a DNA encoding a polypeptide having at least 80% amino acid sequence identity with native sequence DNA19355 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1), wherein said encoded polypeptide activates NF- κ B in a mammalian cell.

15. The nucleic acid molecule of claim 14, wherein said encoded polypeptide has at least 90% amino acid sequence identity.

10

16. The nucleic acid molecule of claim 15, wherein said encoded polypeptide has at least 95% amino acid sequence identity.

17. A vector comprising the nucleic acid of any one of claims 14 to 16.

18. The vector of claim 17, operably linked to control sequences recognized by a host cell transformed with the vector.

20

19. A host cell comprising the vector of claim 17.

20. The host cell of claim 19, which is a CHO cell.

25

21. The host cell of claim 19, which is an *E. coli*.

22. The host cell of claim 19, which is a yeast cell.

23. A process for producing DNA19355 polypeptides, comprising culturing the host cell of any one of claims 19 to 22 under conditions suitable for expression of the polypeptide and recovering the polypeptide from the cell culture.

35

24. The isolated nucleic acid molecule of claim 3, comprising a DNA encoding a soluble polypeptide having at



- 66 -

least 80% amino acid sequence identity with the extracellular domain sequence of a DNA19355 polypeptide consisting of amino acid residues 52 to 177 of Fig. 1 (SEQ ID NO:1), wherein said encoded soluble polypeptide can
5 bind GITR receptor or stimulate mammalian T cells to secrete TNF- α .

25. The nucleic acid molecule of claim 24, wherein said encoded polypeptide has at least 90% amino acid
10 sequence identity.

26. The nucleic acid molecule of claim 24, wherein said encoded polypeptide has at least 95% amino acid
15 sequence identity.

27. An isolated nucleic acid molecule comprising DNA encoding

(a) a DNA19355 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1), or

20 (b) a fragment of (a) which can induce apoptosis in a mammalian cell, activate NF- κ B in a mammalian cell, bind to GITR receptor or stimulate mammalian T cells to secrete TNF- α .

28. An isolated nucleic acid molecule comprising DNA encoding a DNA19355 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1).

29. A vector comprising the nucleic acid of any one
30 of claims 24 to 28.

30. The vector of claim 29, operably linked to control sequences recognized by a host cell transformed with the vector.

31. A host cell comprising the vector of claim 29.



- 67 -

32. The host cell of claim 31, which is a CHO cell.
33. The host cell of claim 32, which is an *E. coli*.
- 5 34. The host cell of claim 33, which is a yeast cell.
35. A process for producing DNA19355 polypeptides, comprising culturing the host cell of any one of claims 31 to 34 under conditions suitable for expression of the polypeptide and recovering the polypeptide from the cell culture.
- 10 36. An isolated nucleic acid molecule comprising the cDNA insert of the vector deposited as ATCC accession number 209466.
- 15 37. Isolated DNA19355 polypeptide comprising amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1).
- 20 38. A polypeptide having at least 96% amino acid sequence identity with native sequence DNA19355 polypeptide consisting of amino acid residues 1 to 177 of Fig. 1 (SEQ ID NO:1), wherein said polypeptide has the ability to
- 25 1) induce apoptosis in a mammalian cell,
ii) activate NF-KB in a mammalian cell,
iii) bind GITR receptor, or
iv) stimulate mammalian T cells to secrete TNF-
α.
- 30 39. A method of inducing apoptosis in mammalian cancer cells, comprising exposing mammalian cancer cells to an effective amount of the polypeptide of claim 38.
- 35 40. A method of stimulating a proinflammatory response in mammalian cells, comprising exposing mammalian cells to an effective amount of the polypeptide of claim



- 68 -

38.

41. The method of claim 40, wherein said mammalian cells are T cells.

5

42. An isolated nucleic acid molecule comprising the cDNA insert of the vector deposited as ATCC 209466.

43. A polypeptide according to claim 38, when used in a method according to any one of claims 39 to 41.

10

44. Use of a polypeptide according to claim 38 for the manufacture of a medicament for inducing apoptosis in mammalian cancer cells.

15

45. Use of a polypeptide according to claim 38 for the manufacture of a medicament for stimulating a proinflammatory response in mammalian cells.

46. An isolated nucleic acid according to any one of claims 1, 3, 14, 24, 27, 28 and 36, substantially as herein described with reference to the examples and drawings.

20

47. An isolated polypeptide according to claim 38, substantially as herein described with reference to the examples and drawings.

25

Dated this 28th day of February 2003

30 GENENTECH, INC.

By their Patent Attorneys

GRIFFITH HACK

Fellows Institute of Patent and

Trade Mark Attorneys of Australia



1 / 11

```

1 CAGCTCTCAT TTCTCCAAA ATGTGTTTGA GCCACTTGGA AAATATGCCT
  GTCGAGAGTA AAGAGGTTTT TACACAACT CGGTGAACCT TTTATACGGA
1 MetCysLeuS erHisLeuGl uAsnMetPro

51 TTAAGCCATT CAAGAAGTCA AGGAGCTCAG AGATCATCCT GGAAGCTGTG
  AATTCGGTAA GTTCTTGAGT TCCTCGAGTC TCTAGTAGGA CCTTCGACAC
  LeuSerHisS erArgThrGl nGlyAlaGln ArgSerSerT rpLysLeuTrp

101 GCTCTTTTGC TCAATAGTTA TGTGCTATT TCTTTGCTCC TTCAGTTGGC
  CGAGAAAACG AGTTATCAAT ACAACGATAA AGAAACGAGG AAGTCAACCG
28 LeuPheCys SerIleValM etLeuLeuPh eLeuCysSer PheSerTrpL

151 TAATCTTTAT TTTTCTCCAA TTAGAGACTG CTAAGGAGCC CTGTATGGCT
  ATTAGAATA AAAAGAGGTT AATCTCTGAC GATTCTCGG GACATACCGA
  euIlePheIl ePheLeuGln LeuGluThrA laLysGluPr oCysMetAla

201 AAGTTTGGAC CATTACCCTC AAAATGGCAA ATGGCATCTT CTGAACCTCC
  TTCAAACCTG GTAATGGGAG TTTTACCGTT TACCGTAGAA GACTTGGAGG
61 LysPheGlyP roLeuProSe rLysTrpGln MetAlaSerS erGluProPr

251 TTGCGTGAAT AAGGTGTCTG ACTGGAAGCT GGAGATACTT CAGAATGGCT
  AACGCACTTA TTCCACAGAC TGACCTTCGA CCTCTATGAA GTCTTACCGA
  oCysValAsn LysValSera spTrpLysLe uGluIleLeu GlnAsnGlyLeu

301 TATATTTAAT TTATGGCCAA GTGGCTCCCA ATGCAAACTA CAATGATGTA
  ATATAAATTA AATACCGGTT CACCGAGGGT TACGTTTGAT GTTACTACAT
95 TyrLeuIl eTyrGlyGln ValAlaProA snAlaAsnTy rAsnAspVal

351 GCTCCTTTTG AGGTGCGGCT GTATAAAAAC AAAGACATGA TACAAACTCT
  CGAGGAAAAC TCCACGCCGA CATATTTTGG TTTCTGTACT ATGTTTGAGA
  AlaProPheG luValArgLe uTyrLysAsn LysAspMetI leGlnThrLeu

401 AACAAACAAA TCTAAAATCC AAAATGTAGG AGGGACTTAT GAATTGCATG
  TTGTTTGTTT AGATTTTAGG TTTTACATCC TCCCTGAATA CTTAACGTAC
128 ThrAsnLys SerLysIleG lnAsnValGl yGlyThrTyr GluLeuHisV

451 TTGGGGACAC CATAGACTTG ATATTCAACT CTGAGCATCA GGTCTAAAA
  AACCCCTGTG GTATCTGAAC TATAAGTTGA GACTCGTAGT CCAAGATTTT
  alGlyAspTh rIleAspLeu IlePheAsnS erGluHisGl nValLeuLys

501 AATAATACAT ACTGGGGTAT CATTTTACTA GCAAATCCCC AATTCATCTC
  TTATTATGTA TGACCCCATTA GTAAAATGAT CGTTTAGGGG TTAAGTAGAG
161 AsnAsnThrT yrTrpGlyIl eIleLeuLeu AlaAsnProG lnPheIleSe

551 CTAGAGACTT GATTGATCTT CCTCATTCCTT TTCAGCACAT GTAGAGGTGC
  GATCTCTGAA CTAACCTAGA GGAGTAAGGG AAGTCGTGTA CATCTCCACG
  rAM*

```

FIG. 1A

601 CAGTGGGTGG ATTGGAGGGA GAAGATATTC AATTCTAGA GTTGTCTGT
GTCACCCACC TAACCTCCCT CTTCTATAAG TTAAAGATCT CAAACAGACA

651 CTACAAAAT CAACACAAAC AGAACTCCTC TGCACGTGAA TTTTCATCTA
GATGTTTTTA GTTGTGTTG TCTTGAGGAG ACGTGCACCT AAAAGTAGAT

701 TCATGCCTAT CTGAAAGAGA CTCAGGGGAA GAGCCAAAGA CTTTGGTGTG
AGTACGGATA GACTTTCTCT GAGTCCCCTT CTCGGTTTCT GAAAACCAAC

751 GATCTGCAGA AATACTTCAT TAATCCATGA TAAAACAAAT ATGGATGACA
CTAGACGTCT TTATGAAGTA ATTAGGTACT ATTTGTTTTA TACCTACTGT

801 GAGGACATGT GCTTTTCAA GAATCTTTAT CTAATTCCTG AATTCATGAG
CTCCTGTACA CGAAAAGTTT CTTAGAAATA GATTAAGAAC TTAAGTACTC

851 TGGAAAAATG GAGTCTATT CCCATGGAAG ATTTACCTGG TATGCAAAAA
ACCTTTTAC CTCAGATAA GGGTACCTTC TAAATGGACC ATACGTTTTT

901 GGATCTGGGG CAGTAGCCTG GCTTTGTTCT CATATTCCTG GGCTGCTGTA
CCTAGACCCC GTCATCGGAC CGAAACAAGA GTATAAGAAC CCGACGACAT

951 ATTCATTCTT CTCATACTCC CATCTTCTGA GACCCTCCCA ATAAAAAGTA
TAAGTAAGAA GAGTATGAGG GTAGAAGACT CTGGGAGGGT TATTTTTCAT

1001 GACTGATAGG ATGGCCACAG ATATGCCTAC CATACCCTAC TTTAGATATG
CTGACTATCC TACCGGTGTC TATACGGATG GTATGGGATG AAATCTATAC

1051 GTGGTGTTAG AAGATAAAGA ACAATCTGAG AACTATTGGA ATAGAGGTAC
CACCACAATC TTCTATTTCT TGTTAGACTC TTGATAACCT TATCTCCATG

1101 AAGTGGCATA AAATGGAATG TACGCTATCT GGAATTTTCT CTTGGTTTTA
TTCACCGTAT TTTACCTTAC ATGCGATAGA CCTTTAAAGA GAACCAAAT

1151 TCTTCCTCAG GATGCAGGT GCTTTAAAAA GCCTTATCAA AGGAGTCATT
AGAAGGAGTC CTACGTCCCA CGAAATTTTT CGGAATAGTT TCCTCAGTAA

1201 CCGAACCTC ACGTAGAGCT TTGTGAGACC TTAATGTTGG TGTGTGTGTC
GGCTTGGGAG TGCATCTCGA AACACTCTGG AATGACAACC ACACACACAG

1251 TAAACATTGC TAATTGTAA GAAAGAGTAA CCATTAGTAA TCATTAGGTT
ATTTGTAACG ATTAACATTT CTTTCTCATT GGTATCATT AGTAATCCAA

FIG. 1B

3 / 11

1301 TAACCCAGAG ATGGTATTAT CATTACTGGA TTATGTCATG TAATGATTTA
ATTGGGGTCT TACCATAATA GTAATGACCT AATACAGTAC ATTACTAAAT

1351 GTATTTTATAG CTAGCTTTCC ACAGTTTGCA AAGTGCTTTC GTAAAAACAGT
CATAAAATC GATCGAAAGG TGTCAAACGT TTCACGAAAG CATTTTGTCA

1401 TAGCAATTCT ATGAAGTTAA TTGGGCAGGC ATTTGGGGGA AAATTTTAGT
ATCGTTAAGA TACTTCAATT AACCGTCCG TAAACCCCTT TTTAAATCA

1451 GATGAGAATG TGATAGCATA GCATAGCCAA CTTTCCTCAA CTCATAGGAC
CTACTCTTAC ACTATCGTAT CGTATCGGTT GAAAGGAGTT GAGTATCCTG

1501 AAGTGACTAC AAGAGGCAAT GGGTAGTCCC CTGCATTGCA CTGTCTCAGC
TTCAGTGATG TTCTCCGTTA CCCATCAGGG GACGTAACGT GACAGAGTCG

1551 TTTAGAATTG TTATTTCTGC TATCGTGTTA TAAGACTCTA AAACCTAGCG
AAATCTTAAC AATAAGACG ATAGCACAAT ATTCTGAGAT TTTGAATCGC

1601 AATTCACTTT TCAGGAAGCA TATTCCCCTT TAGCCCAAGG TGAGCAGAGT
TTAAGTGAAA AGTCCTTCGT ATAAGGGGAA ATCGGGTTCC ACTCGTCTCA

1651 GAAGCTACAA CAGATCTTTC CTTTACCAGC ACACTTTTTT TTTTTTTTCC
CTTCGATGTT GTCTAGAAAG GAAATGGTCG TGTGAAAAAA AAAAAAAGG

1701 TGCCTGAATC AGGGAGATCC AGGATGCTGT TCAGGCCAAA TCCCAACCAA
ACGGACTTAG TCCCTCTAGG TCCTACGACA AGTCCGGTTT AGGGTTGGTT

1751 ATTCCCCTTT TCACTTTGCA GGGCCCATCT TAGTCAAATG TGCTAACTTC
TAAGGGGAAA AGTGAAACGT CCCGGGTAGA ATCAGTTTAC ACGATTGAAG

1801 TAAAATAATA AATAGCACTA ATTCAAAATT TTTGGAATCT TAAATTAGCT
ATTTTATTAT TTATCGTGAT TAAGTTTAA AAACCTAGA ATTTAATCGA

1851 ACTTGCNGGT TGCTTGTTGA AAGGNATATA ATGATTACAT TGTAAACAAA
TGAACGNCCA ACGAACAACT TTCCNTATAT TACTAATGTA ACATTTGTTT

1901 TTTAAATAT TTATGGATAT TTGTGAAAAG CTGCATTATG TTAAATAATA
AAATTTTATA AATACCTATA AACACTTTTC GACGTAATAC AATTTATTAT

1951 TTACATGTAA AGCT
AATGTACATT TCGA

FIG. 1C

4/11

		A		A'		B'	
DNA	19355	52	ETAKEPCMAKFG	-----	PLPSK	-----	WQMASSEP-PCVNKVSDWK--
TNF- α	84	PSDK-PVAHVVA	-----	NQAEQ-QLQ	-----	WLNRR-ANALLANGVELRDNQ	
Apo2L	119	GPQR-VAAHITGTRGRSNTLSSPNSKNEKALGRKINSWESSRSGHFLSNLH-LRNGE					
CD95L	142	E-LR-KVAHLTG	-----	KNSRSM-PL	-----	WEDTY-GIVLLS-GVKYKGG	
LT- α	59	STLK-PAAHLIG	-----	DPSKQN-SLL	-----	WRANT-DRAFLQDGFSLSNNS	
		B		C		D	
DNA	19355	86	LEILQNGLYLIYGVAPNAN	-----	YNDVAPFEVRLYKNK-DMIQTLTNK-SKIQN		
TNF- α	124	LVPSEGLYLIYSQVLFKGQCP	-----	STHVLTHITISRIAS	-----	YQTKVNLLSAIKS	
Apo2L	175	LVIHEKGFYIYSQTYFRFQEEIKENTKIDKQMVQYIYKYTSYDPI--LLMKSARNSC					
CD95L	182	LVINETGLYFVYSKVYFRGQSC	-----	NNLPLSHKVYMRNSKY--PQDLVMMEGKMMS			
LT- α	99	LLVPTSGIYFVYSQVVFSGKAYSPKATSSPLYLAHEVQLFSSQYPFHVPLL-SSQKMVY					
		F		G		H	
DNA	19355	136	-----	VGGTYELHVGDITDILFNSEHQVLKQNT-YWGIILLANPQF-IS			
TNF- α	176	PCQRETPEGAEAKPWYEPYIYLGVGQLEKGDRLSAEINRPDYLDFAESGVYFGIIAL					
Apo2L	232	-----	WSKDAEYGLYSIYQCGIFELKENDRIFVSVTNEHLIDMDHEA-SFFGAFLVG				
CD95L	233	-----	YCTTGQMWARSYLGAVFNLTADHLYVNVSELSLVNF-EESQTFPGLYKL				
LT- α	157	-----	PGLQEPWLHSMYHGAFFLTQGDQLSTHTDGIPHLVL-SPSTVFFGAFAL				

FIG. 2

5/11




FIG. 3




FIG. 4

6/11




FIG. 5A-1 FIG. 5A-2 FIG. 5A-3




FIG. 5A-4 FIG. 5A-5 FIG. 5A-6




FIG. 5B

7/11




FIG. 6




FIG. 8

hG1TR 1 MAQHGAHGA FRATCGALALCAATSLGQRP-TGGFGCGPGRLLLIGITDARQ CRD1
 mG1TR 1 -----MGNWAMLYGVSMLCVLILGQPSVVEEPGCGPGKVQNGSGENNTRO
 hG1TR 50 ORVHTTRCCRDYFGECCSEWDCMEVQPFPHCGDRCCTTCRRHPQPPGQG CRD2
 mG1TR 45 CSLXA-----EGKEDSPKERCICVTPEYHCGDFCKLCKHVECPGGR
 hG1TR 100 VQSQCKFSFGPQCLDCASTFGCGHEHCKPWNDCITQFGFLTVPGNKTH CRD3
 mG1TR 88 MESQGDIVFGERVACAMGTFSAGRDGHCRRLATNESQFGFLTVPGNKTHTM
 hG1TR 150 NAVQVFGSPFAEPLGALTWVILAVAAQVLLLTSAQLGLHIWQLRSQCMMP
 mG1TR 138 NAVCIEEPLDETEYEHITVIFLMAACIFFLTIVQLGHHINOLRQHMCE
 hG1TR 200 RETQLLLEVPFSTEDARSCQFPEERGERSAEEKQRICQDLNV
 mG1TR 188 RETQPPFAEVQISAEIDACSFQFPEERGERQ-TEEXCHLGGRMP

FIG. 7

9/11




FIG. 9A




FIG. 9B




FIG. 10




FIG. 11

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.